

# Interkokask Concentrate 1kg

*DEFRA approved Broad Spectrum Disinfectant*

- A disinfectant suitable for using where keeping poultry; for coop and equipment
- Effective against bacteria, viruses and fungi, red mites and mite eggs, worms and worm eggs, coccidial oocysts and Cryptosporidium

Interkokask has a special formula including chlorocresol which is a lipid-solvent that breaks through the protective triple outer protein/lipid/chitin layers of worm eggs and coccidial oocysts. Interkokask removes immature parasites, adults and larvae.

## Active ingredients

Chlorocresol

## Instructions for use

Note: wear protective clothing; gloves and overalls and read label before use.

1. Remove all animals and dry clean your housing thoroughly.
2. Wash the surfaces using a power washer or brush and water hose.
3. Let the surface dry
4. Apply the diluted product (as advised on the pack) using a hand sprayer or back pack sprayer.
5. Coccidia/Worms/Cryptosporidium use at 3%: Treat the floors and walls up to a minimum of 2 metres along with feeders and drinkers.
6. Mites, lice, use at 3%
7. Viruses and bacteria use at 1%: Treat the whole house.
8. The buildings and equipment should be treated for 2-24 hrs as shown in the 'treatment times' chart below.
9. After the building is dry, re-equip and restock with your birds or other animals.
10. Important: After use, rinse all tools and plastic parts (e.g. High pressure tools, hose pipes) thoroughly with a diluted detergent solution (washing up liquid is a detergent).
11. Wash hands after use.



When using/working with concentrated disinfectants do not eat, drink or smoke. Avoid contact with eyes skin and clothing. Wear protective clothing.

Safe for use on vehicles, machinery and equipment: at recommended dilutions.

Works at low temperatures: DEFRA tested at 4°C.

### Spray Application

- Knapsack spraying equipment: Ideally use a wide angle fan spray nozzle for best coverage.
- Apply a minimum of 100ml solution per square metre on concrete and other smooth surfaces, wooden or porous surfaces should be saturated and may require up to 400ml solution m<sup>2</sup>.

Pack Size: 1 kg (as a liquid).

### 1 kg Interkokask coverage per dilution rate

0.50%	Maximum treatable area approx.	1,920m <sup>2</sup> - 480m <sup>2</sup> *
1.00%	Maximum treatable area approx.	960m <sup>2</sup> -240m <sup>2</sup> *
2.00%	Maximum treatable area approx.	480m <sup>2</sup> -120m <sup>2</sup> *
3.00%	Maximum treatable area approx.	320m <sup>2</sup> - 80m <sup>2</sup> *
4.00%	Maximum treatable area approx.	240m <sup>2</sup> - 60m <sup>2</sup> *
*depending on application rate		

### Treatment Times Chart

UK DEFRA Approvals:			DVG (German Veterinary Association) approved rates:		
General Orders	2.00%	30mins	Disinfection of red mite and red mite eggs	3.00%	24hrs
Poultry Orders	2.00%	30mins	Disinfection of coccidial oocysts	4.00%	2hrs
			Disinfection of cryptosporidium	2.00%	2hrs
			Disinfection of worms and worm eggs	2.00%	2hrs
			Disinfection of enveloped viruses	1.00%	2hrs
European Approvals DIN/EN 14349 bacteria:			DIN/EN 1657 fungi:		

Bacteria (light soiling)	0.50%	30mins	Fungi (light soiling)	2.00%	30mins
Bacteria (heavy soiling)	0.50%	30mins			
DIN/EN 14204 mycobacterium:			Footbaths		
Mycobacterium (Light soiling)	0.50%	60mins	Bacteria/viruses/fungi (coccidial oocysts)	2.00% (3.00%)	Change solution weekly or more often if heavily soiled.
Mycobacterium (Heavy soiling)	1.00%	60mins			

Note: Interkokask is not known to be corrosive at recommended ready-to-use dilutions.

Full instructions are on the pack.